4% Chlorhexidine Solution



(1) Always Read Product Label and Material Safety Data Sheet before use. For Professional use only.

- Contains 4% w/w Chlorhexidine and is the standard solution for surgical hand disinfection.
- Proven effective against bacteria and yeast micro-organisms.

For Surgical Hand Disinfection (EN12791)

Instructions for use:

- Apply 5 ml on hands and scrub for 60 seconds, wash and rinse under running tap water at 20°C for 15 seconds.
- 2. Apply 5 ml on hands and scrub for 2 minutes. Clean fingernails with brush.
- 3. Final rinse for 10 seconds, then dry hands for 5 seconds.

Tested to EN standards PT1 EN12791, EN13624, EN13737

4% Chlorhexidine Solution Test Portfolio

EN 12791

Surgical Hand Disinfection – 3 min Pseudomonas aeruginosa – 3 min Escherichia coli – 3 min Staphylococcus aureus – 3 min Enterococcus hirae – 3 min Candida albicans – 3 min

EN 13624

Yeasticidal – 3 min Candida albicans – 3 min

EN13727

Bactericidal – 60 secs Pseudomonas aeruginosa – 60 secs Escherichia coli – 60 secs Staphylococcus aureus – 60 secs Enterococcus hirae – 60 secs

g9chemicals.com





Your only solution